

Semiclassical Approximations and Predictability in Ocean Acoustics

Michael G. Brown
Division of Applied Marine Physics
Rosenstiel School of Marine and Atmospheric Science
University of Miami
4600 Rickenbacker Causeway
Miami FL 33149-1098
phone (305)361-4640 fax (305)361-4701 email mbrown@rsmas.miami.edu

Award Number N000149710046
<http://www.rsmas.miami.edu/divs/amp.html>

Thrust Category: Long-Range Propagation

LONG-TERM GOALS

Our long-term goal is to improve our understanding of the physics of the forward scattering of underwater sound. Of particular interest is understanding limitations on the predictability of underwater sound fields. Improving our understanding of the mechanism(s) that contribute to the loss of predictability will lead to improved predictive models.

OBJECTIVES

The scientific objective of this work is to understand limitations on the predictability of acoustic wavefields in realistic (range-dependent) ocean environments. Both full wave and semiclassical (ray-based) predictive models are of interest. Both cw and broadband wavefields are of interest. We seek to both develop improved predictive acoustic models and to understand their limitations.

APPROACH

Using semiclassical methods, ideas relating to ray chaos provide a framework for studying predictability. Wavefield behavior in the ray limit is explored numerically and, whenever possible, analytically. The extent to which limitations on predictability carry over to finite frequency wavefields are then explored numerically, generally with parabolic-equation-based models. Both deterministic and stochastic models are of interest. Comparisons with data are made whenever possible.

Several tools have recently been developed and/or extended for use in our work: Tappert's c_o -insensitive PE model (Tappert et al., 1995); the MaCh1 algorithm (Brown, 1994) for finite frequency ray-based wavefield predictions; a stochastic ray model (Brown and Viechnicki, 1998) to model scattering by internal waves; and an efficient algorithm to generate realistic internal-wave-induced sound speed perturbation fields (Colosi and Brown, 1998). This work is being done in loose collaboration with the following individuals: F. Tappert (Univ. Miami; parabolic wave equations, waves in random media, ray chaos); M. Wolfson (Washington State Univ.; numerical modelling, waves in random media, ray chaos); S. Tomsovic (Washington State Univ.; classical and quantum chaos, semiclassical breakdown); G. Zaslavsky (Courant Institute, NYU; chaos in dynamical systems,

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2000		2. REPORT TYPE		3. DATES COVERED 00-00-2000 to 00-00-2000	
4. TITLE AND SUBTITLE Semiclassical Approximations and Predictability in Ocean Acoustics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Division of Applied Marine Physics,,Rosenstiel School of Marine and Atmospheric Science,,University of Miami,4600 Rickenbacker Causeway,,Miami,,FL, 33149				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Our long-term goal is to improve our understanding of the physics of the forward scattering of underwater sound. Of particular interest is understanding limitations on the predictability of underwater sound fields. Improving our understanding of the mechanism(s) that contribute to the loss of predictability will lead to improved predictive models.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

stochastic methods, quantum chaos); and J. Colosi (WHOI; ocean internal waves and long-range underwater sound propagation).

WORK COMPLETED

Recent accomplishments include: 1) development and numerical implementation of stochastic ray theory (Brown and Viechnicki, 1998) to describe and model the forward scattering of underwater sound by internal waves; 2) development of a new efficient, accurate and robust algorithm to compute internal-wave-induced sound speed perturbation fields (Colosi and Brown, 1998); 3) incorporation of this internal wave model into both ray and full wave propagation models; 4) demonstration (Brown, 1999) that, in the ray limit, islands of stability in phase space exist when the environment is almost stratified and has nonperiodic range-dependence; and 5) construction (Brown, 1999) of the solution, in the form of a Maslov integral, of a large class of problems involving wave motion in inhomogeneous moving media.

RESULTS

Stochastic ray theory and the new technique to compute internal-wave-induced sound speed perturbation fields are important new tools which can be used to address fundamental questions relating to the predictability of long-range sound propagation in deep ocean environments. The work relating to the structure of phase space provides critically important insight into -- and a mathematical framework for the study of -- sound propagation in environments with nontrivial range-dependence. Our (with M. Wolfson and S. Tomsovic) semiclassical breakdown studies show that, even in the presence of chaotic ray motion, semiclassical wavefield representations accurately predict many important features of underwater acoustic wavefields. Recent work on analysis of the AET long-range propagation data indicates that path stability is controlled to a large extent by the background sound speed structure.

IMPACT/APPLICATIONS

Our work gives insight into the limitations on the predictability of underwater sound fields. This is an important basic science issue which impacts all systems applications which require accurate predictions of underwater sound fields.

TRANSITIONS

The PI collaborates informally with the investigators listed above, ATOC investigators, and others. This includes sharing both ideas and software. It is not known whether any software produced by the PI has been used to address any applied Navy problems.

RELATED PROJECTS

This work is closely related to the ATOC project and the ONR-funded work being performed by P. Worcester (SIO), J. Colosi (WHOI), M. Wolfson (WSU), J. Spiesberger (UPenn), S. Tomsovic (WSU), G. Zaslavsky (CIMS/NYU) and F. Tappert (UMiami). All of these projects are concerned with aspects of long-range propagation.

REFERENCES

Brown, M. G. (1994) A Maslov-Chapman wavefield representation for wide-angle one-way propagation, *Geophys. J. Intl.* 116, 523-526.

Brown, M. G. (1998) Phase space structure and fractal trajectories in $1\frac{1}{2}$ degree of freedom Hamiltonian systems whose time dependence is quasiperiodic, *Nonlinear Processes in Geophys.* 5, 69-74.

Brown, M. G., and J. Viechnicki (1998) Stochastic ray theory for long-range sound propagation in deep ocean environments, *J. Acoust. Soc. Am.* 104, 2090-2104.

Colosi, J. A., and M. G. Brown (1998) Efficient numerical simulation of stochastic internal-wave-induced sound speed perturbation fields, *J. Acoust. Soc. Am.* 103, 2232-2235.

Tappert, F. D., Spiesberger, J. L., and L. Boden (1995) New full-wave approximation for ocean acoustic travel time predictions, *J. Acoust. Soc. Am.* 97, 2771-2782.

PUBLICATIONS

Brown, M. G. (1999) The Maslov integral representation of slowly-varying dispersive wavetrains in inhomogeneous moving media, *Wave Motion* 32, 247-266.

Brown, M. G., J. A. Colosi, A. L. Virovlyansky, G. M. Zaslavsky, S. Tomsovic and M. A. Wolfson (2000) Ray dynamics in the AET experiment, in preparation.

Brown, M. G., and M. A. Wolfson (2000) A numerical investigation of semiclassical breakdown in the presence of ray chaos, in preparation.